

Original Article
Obstetrics & Gynecology



Women with Endometriosis, Especially Those Who Conceived with Assisted Reproductive Technology, Have Increased Risk of Placenta Previa: Meta-analyses

Hyeji Jeon , Jiwon Min , Deok Kyeong Kim , Heekyung Seo ,
Sunkyung Kim , and Yun-Sook Kim

Department of Obstetrics and Gynecology, Soonchunhyang University College of Medicine, Soonchunhyang University Cheonan Hospital, Cheonan, Korea



Received: Mar 8, 2018
Accepted: Jul 18, 2018

Address for Correspondence:

Yun-Sook Kim, MD, PhD

Department of Obstetrics and Gynecology,
Soonchunhyang University, College of Medicine,
Soonchunhyang University Cheonan Hospital,
31 Soonchunhyang 6-gil, Dongnam-gu,
Cheonan, 31151, Republic of Korea.
E-mail: drsook@schmc.ac.kr

© 2018 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Hyeji Jeon
<https://orcid.org/0000-0001-9171-8349>
Jiwon Min
<https://orcid.org/0000-0003-0192-5411>
Deok Kyeong Kim
<https://orcid.org/0000-0001-5300-5524>
Heekyung Seo
<https://orcid.org/0000-0003-4886-2374>
Sunkyung Kim
<https://orcid.org/0000-0002-6161-3535>
Yun-Sook Kim
<https://orcid.org/0000-0001-8427-4006>

ABSTRACT

Background: Many women with endometriosis have become pregnant through assisted reproductive technology (ART), and have often experienced placenta previa (PP) during pregnancy. The objective of this study was to assess the association between women with endometriosis, especially those who conceived with ART, and the risk of PP.

Methods: Two reviewers independently determined studies that were considered suitable for meta-analyses published in various medicine-related databases from March 1, 2004 through July 31, 2017 without language restrictions. Eight studies met the inclusion criteria, with a combined sample size of 21,930 women. Of these 21,930 pregnancies, 6,256 had endometriosis (endometriosis) and 15,674 had no endometriosis. Four of these studies included 8,161 women who conceived with ART, 1,640 of whom had endometriosis (endometriosis + ART), and 6,521 of whom did not have endometriosis. Meta-analyses were estimated with odds ratios (ORs) and 95% confidence intervals (CIs) using random effect analysis according to heterogeneity of studies.

Results: These meta-analyses showed women with endometriosis (endometriosis) have an increased risk of PP (OR, 4.038; 95% CI, 2.291–7.116; $P = 0.000$). These results showed women who conceived with ART (endometriosis + ART), have a substantially increased risk of PP (OR, 5.543; 95% CI, 1.659–18.523; $P = 0.005$).

Conclusion: These meta-analyses demonstrate women with endometriosis have an increased risk of PP.

Keywords: Endometriosis; Placenta Previa; Meta-analysis; Assisted Reproductive Technology

INTRODUCTION

Placenta previa (PP) is one of the diseases in which the placenta covers the internal os of the cervix and causes obstetric hemorrhagic complications during pregnancy and cesarean section.¹ The incidence of this disease is about 0.3%–0.5%, and it is an important disease because it increases maternal and perinatal morbidity and mortality from massive bleeding.² The risk factors for PP include a previous history of cesarean section, increasing

Disclosure

Within the past 12 months, the authors had no financial relationships with any biotechnology manufacturers, pharmaceutical companies, or other commercial entities that have an interest in the subject matter or materials discussed in this manuscript.

Author Contributions

Conceptualization: Kim YS. Data curation: Jeon H, Min J. Formal analysis: Seo H, Kim S. Methodology: Min J, Kim DK. Software: Kim YS, Kim S. Validation: Kim YS, Jeon H. Investigation: Kim DK, Seo H. Writing - original draft: Jeon H. Writing - review & editing: Kim YS.

maternal age, multiple pregnancy, and smoking, and recent reports have indicated that assisted reproductive technology (ART) is also a cause.³⁻⁵ However, the pathogenesis of PP has not been clearly understood until recently. Most cases of PP should be delivered by caesarean section, not by vaginal delivery. Endometriosis is a chronic reproductive disease characterized by the presence of endometrial glands outside the uterus, mainly on the ovaries and peritoneum. The prevalence of endometriosis in women of reproductive age is about 6%–10%.⁶ Endometriosis causes two main symptoms, pain and infertility. It is well accepted that endometrium of women with endometriosis is abnormal. Additionally, these abnormal endometrium may be the causes of decidual impairment and abnormal placentation during pregnancy.⁷ Endometriosis has been linked with increased peritoneal inflammation with a higher concentration of cytokines, angiogenic factors, and growth factors. Other harmful factors include poorer oocyte quality, progesterone receptor resistance, molecular and functional abnormalities in eutopic endometrium, and anatomical distortion of uterine posterior walls, fallopian tubes and ovaries in women with endometriosis.^{8,9} So, all of these processes may have negative impacts on pregnancy outcomes and fertility. However, studies on the relationship between PP and endometriosis have led to conflicting results, with some studies reporting a significantly increased risk of PP,¹⁰⁻¹⁴ and others showing no statistical significance, though in small sample sizes.¹⁵ Many women with endometriosis tend to have conceived with ART, though it is unclear whether the causes of the increase in PP are due to endometriosis, ART, or both.^{16,17} The objective of our study was to examine whether pregnant women with endometriosis are associated with an increased risk of PP.

METHODS

We developed a search strategy to use Medical Subject Heading (MeSH) terms and free key words and text words related to “endometriosis”, “pregnancy”, “adverse pregnancy outcomes”, “bad pregnancy outcomes”, “placenta previa”, “ART”, “in vitro fertilization (IVF)”, “intracytoplasmic sperm injection (ICSI)”, and “placental problems”.

We searched these terms using the PubMed MEDLINE database, the Korea education and research information service (KERIS), Scopus, and Google Scholar from March 1, 2004 through July 31, 2017 without language restrictions. The inclusion studies included a prospective cohort study, a retrospective cohort study, a retrospective case control study, a large national population-based cohort study, a retrospective secondary analysis, and double blinded, multicentric, observational and cohort studies, and a placebo-controlled, randomized clinical trial on the relations between endometriosis and pregnancy prognosis, while publications in abstract form alone were excluded. Data abstraction was completed by two independent investigators. Each investigator independently abstracted data from each study and analyzed these data separately.

Inclusion and exclusion criteria

The endometriosis group had pelvic endometriosis confirmed histologically and visually during the surgical procedure, or in two or more repetitive ultrasound examinations or magnetic resonance imaging (MRI) scans before pregnancy or after the operation. All patients who were clinically suspected to have pelvic endometriosis but with an absence of imaging or surgical confirmation were excluded from the endometriosis group. In this study, ART that was conducted by conventional IVF or ICSI was included. The gamete intrafallopian transfer (GIFT) women were excluded. The unexposed group included women who did not

have a previous surgical or clinical diagnosis of endometriosis, and who did not have any ultrasonographic signs of endometriosis. Women with malignancies, autoimmune diseases, endocrine diseases, and cardiovascular diseases were also excluded. Donor oocyte and embryo recipients were excluded. Miscarriages and terminations of pregnancy at < 20 weeks gestation were not registered as births. Late termination \geq 20 weeks gestation and neonatal death (within 28 days of birth) were included.

Definitions

PP is defined as placental tissue completely obstructing internal os or within 20 mm. PP was confirmed at delivery. Confirmation of the diagnosis of PP was based on transvaginal ultrasonography with empty bladder performed prior to delivery. If the PP was not confirmed prior to delivery and there was a sudden unexpected birth, the diagnosis was confirmed at the time of delivery.

Statistical analysis

Differences were reviewed, and further resolved by a common review of the entire data set. Data abstracted included the number of study patients, the number of patients in endometriosis groups and control groups, and adverse pregnancy outcomes. When possible, authors of included trials were contacted for missing data. The risk of bias in each included study was assessed using the criteria outlined in the Risk of Bias Assessment Tool for Non-randomized studies (RoBANS). The following seven domains related to risk of bias were assessed in each included trial since there is evidence that these issues are associated with biased estimates of treatment effect: 1) random sequence generation, 2) allocation concealment, 3) blinding of participants and personnel, 4) blinding of outcome assessment, 5) incomplete outcome data, 6) selective reporting, and 7) other bias. The judgments of review authors were categorized as having a low, high, or unclear risk of bias.

Meta-analyses were performed with random effects models according to the heterogeneity of studies and we attempted to estimate the mean of a distribution of true effects. The completed analyses were then compared, and any differences were resolved with a review of the entire data set and independent analysis. Statistical heterogeneity between the studies was assessed using I^2 statistics. Heterogeneity was considered to be statistically significant if the I^2 value was more than 50%. Publication bias was evaluated by inspection of the funnel plot, Duval and Tweedie's trim and fill, and Begg and Mazumdar's rank correlation. Kendall's tau value and Q value were adjusted to see a bias in publications. All effect sizes were calculated through Comprehensive Meta-analysis (CMA) ver.2.0 (Biostat Inc., Englewood, NJ, USA). Meta-analyses were estimated with odds ratio (OR) and 95% confidence interval (CI) using random effect analysis according to the heterogeneity of studies. A P value less than 0.05 was considered to be statistically significant. The meta-analysis was performed following the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) statement.

RESULTS

Fig. 1 shows a flow diagram of how we identified relevant studies. A total of 246 articles were identified by searching the PubMed, the Embase, the Cochrane Library, and the KERIS. We excluded 45 duplicated articles and an additional 182 articles that did not satisfy the selection criteria. The full texts of the remaining 19 articles were reviewed, and 11 additional articles were excluded for reasons shown in **Fig. 1**. The remaining 8 articles were included in the final




Fig. 1. Flow diagram of identification of relevant studies.
KERIS = Korea Education and Research Information Service.

analysis. The summary of characteristics of the enrolled studies included in these meta-analyses are in order. All of their RoBANS were a low and unclear risk of bias, and they were evaluated as good studies in the assessment of risk of bias (**Table 1**).


Meta-analyses were performed with random effects models according to the heterogeneity of eight studies (endometriosis) as we attempted to estimate the mean of a distribution of true effects. According to Kendall's tau without continuity correction, the tau was -0.214 , the Z value for tau was 0.742 , the P value (1-tailed) was 0.228 , and the P value (2-tailed) was 0.457 . Also, as a result of Duval and Tweedie's trim and fill, the point estimate was shown, and the Q value was 45.053 . It was difficult to see if there was a bias in publications because the adjusted values and observed values of the studies were the same. Figure of the funnel plot for assessing publication bias for the risk of PP and endometriosis (endometriosis) is shown; it looks apparent that the funnel plot is symmetric (**Fig. 2**). Data from eight effect sizes from eight studies involving 21,930 patients were enrolled. These meta-analyses showed that women with endometriosis (endometriosis) have an increased risk of PP (OR, 4.038; 95% CI, 2.291–7.116; $P = 0.000$, **Fig. 3**).

Meta-analyses were performed with random effects models according to the heterogeneity of four studies (endometriosis + ART) as we attempted to estimate the mean of a distribution of true effects. According to Kendall's tau without continuity correction, the tau was 0.000 , the Z value for tau was 0.000 , the P value (1-tailed) was 0.500 , and the P value (2-tailed) was 1.000 . Also, as a result of Duval and Tweedie's trim and fill, point estimate was shown, and the Q value was 19.282 . Figure showed funnel plot for assessing publication bias for the risk of PP conceived by ART (endometriosis + ART); it looked apparent that the funnel plot was also symmetric (**Fig. 4**). It was difficult to see if there was a bias in publications. Data from four effect sizes from four studies involving 8,161 patients who conceived with ART were

Table 1. Summary of characteristics of enrolled studies included meta-analyses

Descriptions	Healy et al. ¹⁰	Takemura et al. ¹¹	Lin et al. ¹²	Tatsuya et al. ¹³	Benaglia et al. ¹⁴	Li et al. ¹⁵	Saraswat et al. ¹⁶	Berliac et al. ¹⁷
Study design	Retrospective cohort study	Retrospective cohort study	Retrospective cohort study	Retrospective cohort study	Retrospective case control study	Retrospective cohort study	National population based cohort study	National cohort study
Type of conception	1,265 ART+En+, 5,465 ART+En-	44 ART+En+, 305 ART+En-	249 non-ART+En+, 249 non-ART+En-	92 ART+En+, 512 ART+En-	239 ART+En+, 239 ART+En-	1 ART+En+, 74 non-ART+En+, 300 non-ART+En-	ART not distinguished	ART not distinguished
Site, year	Australia, 1991-2004	Japan, 2004-2008	China, 1995-2013	Japan, 2000-2014	Italy, 2008-2014	China, 2011-2013	Scotland, 1981-2010	Denmark, 1997-2014
Endometriosis surgery before pregnancy (type & time)	NA	44 (type & time: NA)	249 (type & time: NA)	92 (type: adhesiotomy, coagulation, cystectomy, time: NA)	186 (78%) (type: cystectomy, removal of deep peritoneal nodules, time: NA)	75 (type & time: NA)	4,232 (type & time: NA)	60 (type & time: NA)
Percentage of previous cesarean section (%)	NA	0% ART+En+, 2.6% ART+En-	0% non-ART+En+, 0% non-ART+En-	NA	NA	NA	NA	NA
Percentage of nulliparity (%)	NA	100% ART+En+, 97.4% ART+En-	100% non-ART+En+, 100% non-ART+En-	84.8% ART+En+, 77.6% ART+En-	90% ART+En+, 84% ART+En-	100% En+, 58% En-	72% En+, 52% En-	67% En+, 54% En-
Primary outcomes	Obstetric hemorrhage (APH, PP, PA, PPH)	PP	Adverse obstetric outcomes (preterm labor, PP, cesarean section, PIH, SGA)	Adverse obstetric outcomes (preterm labor, PP, SGA)	Preterm birth	Adverse obstetric outcomes (PPH, preterm labor, PP, SGA, GDM)	Adverse obstetric outcomes (miscarriage, ectopic pregnancy, stillbirths, complications PIH, APH, PPH, preterm labor, cesarean section)	The risk of obstetrical & neonatal complications
Diagnostic methods of endometriosis	NA	44 Histopathology on surgery	249 Histologically & visually on surgery	92 Histopathology on laparoscopic surgery	186 Histopathology on surgery, 53 transvaginal sonography	75 Histopathology on laparoscopic surgery	4,232 Surgically (laparoscopy or laparotomy) confirmed	60 Histopathology on surgery
Sample size (No.) (En+/En-)	6,730 (1,265/5,465)	349 (44/305)	498 (249/249)	604 (92/512)	478 (239/239)	375 (75/300)	10,939 (4,232/6,707)	1,957 (60/1,897)
Adjusted RR/ OR (95% CI)	1,700 (1,220-2,404)	13,347 (3,939-45,294)	4,517 (1,233-16,544)	12,095 (3,563-41,061)	4,800 (1,369-16,824)	1,616 (0,307-8,499)	2,240 (1,518-3,306)	5,700 (4,395-7,392)
Risk of bias	Unclear	Low	Low	Low	Low	Low	Unclear	Unclear

NA = not available, No = number of patients, En+ = endometriosis group, En- = control group, RR = risk ratio, OR = odds ratio, CI = confidence interval, ART = assisted reproductive technology, APH = antepartum hemorrhage, PP = placenta previa, PA = placental abruption, PPH = postpartum hemorrhage, PIH = pregnancy induced hypertension, SGA = small for gestational age, GDM = gestational diabetes mellitus.



Duval and Tweedie's trim and fill

	Fixed effects				Random effects			Q value
	Studies trimmed	Point estimate	Lower limit	Upper limit	Point estimate	Lower limit	Upper limit	
Observed values		3.50632	2.94348	4.17680	4.03753	2.29078	7.11620	45.05037
Adjusted values	0	3.50632	2.94348	4.17680	4.03753	2.29078	7.11620	45.05037

Begg and Mazumdar rank correlation


Kendall's S statistic (P-Q)	-6.00000
Kendall's tau without continuity correlation	
Tau	-0.21429
Z value for tau	0.74231
P value (1-tailed)	0.22895
P value (2-tailed)	0.45790
Kendall's tau without continuity correlation	
Tau	-0.17857
Z value for tau	0.61859
P value (1-tailed)	0.26809
P value (2-tailed)	0.53619

Fig. 2. Funnel plot for assessing publication bias for the PP risk in women with endometriosis (endometriosis). PP = placenta previa.

Study or subgroup	Endometriosis (En+)		Comparator (En-)		Weight	Statistics for each study					OR, random, 95% CI
	Events	Total	Events	Total		OR	Lower limit	Upper limit	Z value	P value	
Healy et al. ¹⁰	48	1,265	126	5,465	17.69%	1.700	[1.202	2.404]	3.001	0.003	
Takemura et al. ¹¹	9	44	4	305	10.09%	13.347	[3.939	45.224]	4.162	0.000	
Lin et al. ¹²	13	249	3	249	9.50%	4.517	[1.233	16.544]	2.276	0.023	
Tatsuya et al. ¹³	8	92	4	512	10.07%	12.095	[3.563	41.061]	3.997	0.000	
Benaglia et al. ¹⁴	14	239	3	239	9.83%	4.800	[1.369	16.824]	2.451	0.014	
Li et al. ¹⁵	2	75	5	300	7.22%	1.616	[0.307	8.499]	0.567	0.571	
Saraswat et al. ¹⁶	72	4,232	54	6,707	17.39%	2.240	[1.518	3.306]	4.062	0.000	
Berlac et al. ¹⁷	2	60	8	1,897	18.22%	5.700	[4.395	7.392]	13.123	0.000	
Total (95% CI)		6,256		15,674	100.00%	4.038	[2.291	7.116]	4.827	0.000	
Total Events	168		207								

Heterogeneity: Tau² = 0.82; Chi² = 45.05, df = 7 (P = 0.000); I² = 84%
 Test for overall effect: Z = 4.827 (P < 0.000)

Fig. 3. Forest plot of eight studies. The result of the PP risk in women with endometriosis (endometriosis). PP = placenta previa, OR = odds ratio, CI = confidence interval.



Duval and Tweedie's trim and fill

	Fixed effects				Random effects			Q value
	Studies trimmed	Point estimate	Lower limit	Upper limit	Point estimate	Lower limit	Upper limit	
Observed values		2.35487	1.72447	3.21571	5.54349	1.65903	18.52307	19.28222
Adjusted values	0	2.35487	1.72447	3.21571	5.54349	1.65903	18.52307	19.28222

Begg and Mazumdar rank correlation

Kendall's S statistic (P-Q)	0.00000
Kendall's tau without continuity correlation	
Tau	0.00000
Z value for tau	0.00000
P value (1-tailed)	0.50000
P value (2-tailed)	1.00000
Kendall's tau without continuity correlation	
Tau	0.00000
Z value for tau	0.00000
P value (1-tailed)	0.50000
P value (2-tailed)	1.00000

Fig. 4. Funnel plot for assessing publication bias for the PP risk in women with endometriosis who were conceived with ART (endometriosis + ART). PP = placenta previa, ART = assisted reproductive technology, OR = odds ratio.

Study or subgroup	ART+En+		Comparator (ART+En-)		Weight	Statistics for each study					OR, random, 95% CI
	Events	Total	Events	Total		OR	Lower limit	Upper limit	Z value	P value	
Healy et al. ¹⁰	48	1,265	126	5,465	30.05%	1.700	[1.202	2.404]	3.001	0.003	
Takemura et al. ¹¹	9	44	4	305	23.43%	13.347	[3.939	45.224]	4.162	0.000	
Tatsuya et al. ¹³	8	92	4	512	23.41%	12.095	[3.563	41.061]	3.997	0.000	
Benaglia et al. ¹⁴	14	239	3	239	23.11%	4.800	[1.369	16.824]	2.451	0.014	
Total (95% CI)		1,640		6,521	100.00%	5.543	[1.659	18.523]	2.782	0.005	
Total Events	79		137								

Heterogeneity: Tau² = 0.81; Chi² = 19.28, df = 3 (P = 0.000); I² = 84%
 Test for overall effect: Z = 2.782 (P = 0.005)

Fig. 5. Forest plot of four studies. The result of the PP risk in women with endometriosis who were conceived with ART (endometriosis + ART). PP = placenta previa, ART = assisted reproductive technology, OR = odds ratio, CI = confidence interval.

included. These results showed that women with endometriosis who conceived with ART (endometriosis+ ART) have a substantially increased risk of PP (OR, 5.543; 95% CI, 1.659–18.523; P= 0.005, Fig. 5).

DISCUSSION

Endometriosis is a common and painful disease affecting women of reproductive age. While the underlying pathophysiology is still largely unknown, much advancement has been made in understanding the progression of the disease. A recent meta-analysis by Kim et al.¹⁸ demonstrated that primiparous singleton women with endometriosis at pregnancy have an increased risk of preterm birth (OR, 1.473; 95% CI, 1.216–1.785). In a study comparing the complications of pregnancy and delivery in pregnant women with endometriosis, it is important to distinguish between whether an accurate laparoscopic diagnosis was given before or after pregnancy and whether it was pregnancy through ART such as IVF/ET or ICSI. A retrospective analysis by Jeon et al.¹⁹ found that complete cul-de-sac obliteration is the independent factor of spontaneous pregnancy failure in women with endometriosis following laparoscopic surgery. In the process of embryo implantation in the endometrium, inadequate uterine contractility called 'endometrial waves' observed in the sub-endometrial layer are thought to be the causes of PP.²⁰ Population studies have found annual incidence rates of endometriosis ranging from 0.1–0.3.²¹ This reflects an increase in the diagnosis of endometriosis. According to the study on progesterone resistance in endometriosis conducted by Chae et al.²² in 2016, it may be caused by proinflammatory conditions in the pelvic peritoneal microenvironment. The biochemical materials associated in implantation include local and systemic inflammatory cytokines such as interleukin-6, interleukin-1 β , and tumor necrosis factor- α . Additionally, the levels of prostaglandin E₂, cyclooxygenase (COX)-2 and various cytokines are highly elevated in endometriotic tissue relative to normal endometrium. The hypermethylation of progesterone receptor (PR)-B promoter and PR isoform A are also more expressed than PR-B as well as in endometriosis.²³ According to a retrospective analysis by Yang et al.,²⁴ endometriosis itself has worse IVF parameters regardless of previous cyst enucleation. In 2012, Vercellini et al.²⁵ retrospectively assessed pregnancy outcomes in 419 women who achieved a first spontaneous singleton pregnancy following surgery for endometriosis, and stratified the results obtained by endometriosis localization. In this study, an almost six-fold increase in risk of PP has been found in women with rectovaginal endometriosis compared to all women with ovarian and peritoneal lesions (OR, 5.81; 95% CI, 1.53–22.03). Because a fixed abnormal anatomical uterine position owing to dense pelvic adhesions may theoretically reduce the efficacy of myometrial contraction, particularly in women with rectovaginal endometriosis, the risk of PP increases. In a small case-control study involving women with and without endometriosis matched for parity and ART procedures conducted by Kortelahti et al.,²⁶ no difference in PP incidence was found. However, the study likely had a very limited statistical power, as the number of placental abnormalities reported was very low. Conversely, two retrospective cohort studies conducted by Healy et al.¹⁰ and Kuivasaari-Pirinen et al.²⁷ comparing the pregnancy outcomes of ART singleton pregnancies with those of natural pregnancies found higher rates of PP in the ART groups and in the various subgroups. The limitation of Healy et al.'s study¹⁰ was no information of previous section history, percentage of nulliparity, and diagnostic methods of endometriosis. Although the study was conducted on only ART pregnancies and may cause bias in the final statistics, it was included in this analysis due to a large number of samples. Kuivasaari-Pirinen et al.²⁷ showed that mothers with endometriosis and those in the male factor infertility group have a greater incidence of PP than the reference group. One of the questions of this study is why ART increases the risk of PP. This study is consistent with other studies in that PP is increased in ART pregnancies. The underlying mechanism for this effect is not clear. In assisted fertilization, drugs are utilized to induce multiple follicular development. Fertilization and embryo development take place outside the body, and

embryos enter the uterine cavity through the cervix and vagina by mechanical means. With ART, embryos are placed in the uterine cavity by the transcervical route using a catheter. This procedure may induce uterine contraction, possibly due to the release of prostaglandins after mechanical stimulation of the internal cervical os. It is conceivable that these mechanically induced abnormal uterine contractions could lead to higher frequencies of implantation in the lower uterine segment and thereby increase the risk of PP. And, the normal frequency and amplitude of uterine contractions are altered in women with endometriosis, which may cause abnormal uterine dysperistalsis in endometriotic tissue leading to abnormal embryo implantation resulting in PP.²⁸ These comorbidities in women with endometriosis who conceived with ART may increase the incidence of PP, not by ART itself.

In 2016, Tatsuya et al.¹³ tried to exclude the negative effects of ART on pregnancy by studying only women who conceived with ART. They also retrospectively investigated whether the severity of endometriosis was associated with increased complications of pregnancy conceived with ARTs. Endometriosis was classified according to revised American Society for Reproductive Medicine (rASRM) staging.²⁹ They found that the frequencies of PP was significantly increased in women with laparoscopically diagnosed endometriosis (OR, 12.1; 95% CI, 3.6–41.1).¹³ The association between ovarian or rectovaginal endometriosis and the increased risks of PP involves the relationship between the severity of endometriosis and increased complications of pregnancy. The prevalence of bad pregnancy outcomes in the first pregnancy has been reported to increase compared to the next pregnancy, so this study has a strength in that it examined only nulliparous women.³⁰

We selected studies of women who underwent pre-pregnancy surgery, except for the study of Healy and others in 2010. The study of Healy et al did not show how endometriosis was diagnosed. And, while the study of Benaglia et al.¹⁴ in 2016 found that only 78% of women had surgery before conception, the other six had surgery all before pregnancy. And the percentage of nulliparity was from 52% to 100%. The diagnostic methods of endometriosis were histology except for 53 transvaginal sonography in the study of Benaglia et al.¹⁴ in 2016. Especially, both multiparity and previous cesarean section history have been known as important risk factors of PP. But, most of the studies included in our meta-analysis did not show previous C-section rates. It is assumed that it will be similar to the percentage of nulliparity. These factors may cause risk of bias in our analysis.

We would like to highlight the important findings and the significance of our analyses. The first strength of our study was that we analyzed the most recent studies available. The second strength of our study was that we performed the meta-analyses by separating the ART plus endometriosis group, taking into account that ART may affect PP development. The limitations of our study are that seven out of the eight studies used in the analysis indicated that endometriosis increased the incidence of PP, and the only other study was not significant. Because it is almost certain that PP increases in women with endometriosis, these analyses focused on how much it increases. The “unexposed” cohort can include some women with endometriosis undiagnosed by surgery or through imaging methods. These inclusion criteria thus tend to underestimate the risk of complications in women with endometriosis. Future research should focus on the use of fresh embryos and FETs and the type of infertility (tubal, ovulatory disorder, and male infertility) used in ART, as well as endometriosis localization (rectovaginal lesions, ovarian endometriomas, and peritoneal implants) and comparisons of the severity of endometriosis (classification of as rASRM stage I, II, III, and IV), followed by correction of maternal age and the study period, which will lead to better studies.

In conclusion, our study demonstrates that women with endometriosis have an increased risk of PP. We must increase international effort to understand the etiology and pathogenesis of endometriosis. Also, obstetrics should try to reduce complications in women with endometriosis from the time of conception to the postpartum period.

REFERENCES

- Oyelese Y, Smulian JC. Placenta previa, placenta accreta, and vasa previa. *Obstet Gynecol* 2006;107(4):927-41. [PUBMED](#) | [CROSSREF](#)
- Iyasu S, Saftlas AK, Rowley DL, Koonin LM, Lawson HW, Atrash HK. The epidemiology of placenta previa in the United States, 1979 through 1987. *Am J Obstet Gynecol* 1993;168(5):1424-9. [PUBMED](#) | [CROSSREF](#)
- Ananth CV, Smulian JC, Vintzileos AM. The association of placenta previa with history of cesarean delivery and abortion: a metaanalysis. *Am J Obstet Gynecol* 1997;177(5):1071-8. [PUBMED](#) | [CROSSREF](#)
- Ananth CV, Wilcox AJ, Savitz DA, Bowes WA Jr, Luther ER. Effect of maternal age and parity on the risk of uteroplacental bleeding disorders in pregnancy. *Obstet Gynecol* 1996;88(4 Pt 1):511-6. [PUBMED](#) | [CROSSREF](#)
- Romundstad LB, Romundstad PR, Sunde A, von Düring V, Skjaerven R, Vatten LJ. Increased risk of placenta previa in pregnancies following IVF/ICSI; a comparison of ART and non-ART pregnancies in the same mother. *Hum Reprod* 2006;21(9):2353-8. [PUBMED](#) | [CROSSREF](#)
- Giudice LC, Kao LC. Endometriosis. *Lancet* 2004;364(9447):1789-99. [PUBMED](#) | [CROSSREF](#)
- Buck Louis GM, Hediger ML, Peterson CM, Croughan M, Sundaram R, Stanford J, et al. Incidence of endometriosis by study population and diagnostic method: the ENDO study. *Fertil Steril* 2011;96(2):360-5. [PUBMED](#) | [CROSSREF](#)
- Stratton P, Berkley KJ. Chronic pelvic pain and endometriosis: translational evidence of the relationship and implications. *Hum Reprod Update* 2011;17(3):327-46. [PUBMED](#) | [CROSSREF](#)
- May KE, Villar J, Kirtley S, Kennedy SH, Becker CM. Endometrial alterations in endometriosis: a systematic review of putative biomarkers. *Hum Reprod Update* 2011;17(5):637-53. [PUBMED](#) | [CROSSREF](#)
- Healy DL, Breheny S, Halliday J, Jaques A, Rushford D, Garrett C, et al. Prevalence and risk factors for obstetric haemorrhage in 6730 singleton births after assisted reproductive technology in Victoria Australia. *Hum Reprod* 2010;25(1):265-74. [PUBMED](#) | [CROSSREF](#)
- Takemura Y, Osuga Y, Fujimoto A, Oi N, Tsutsumi R, Koizumi M, et al. Increased risk of placenta previa is associated with endometriosis and tubal factor infertility in assisted reproductive technology pregnancy. *Gynecol Endocrinol* 2013;29(2):113-5. [PUBMED](#) | [CROSSREF](#)
- Lin H, Leng JH, Liu JT, Lang JH. Obstetric outcomes in Chinese women with endometriosis: a retrospective cohort study. *Chin Med J (Engl)* 2015;128(4):455-8. [PUBMED](#) | [CROSSREF](#)
- Fujii T, Wada-Hiraike O, Nagamatsu T, Harada M, Hirata T, Koga K, et al. Assisted reproductive technology pregnancy complications are significantly associated with endometriosis severity before conception: a retrospective cohort study. *Reprod Biol Endocrinol* 2016;14(1):73-7. [PUBMED](#) | [CROSSREF](#)
- Benaglia L, Candotti G, Papaleo E, Pagliardini L, Leonardi M, Reschini M, et al. Pregnancy outcome in women with endometriosis achieving pregnancy with IVF. *Hum Reprod* 2016;31(12):2730-6. [PUBMED](#) | [CROSSREF](#)
- Li H, Zhu HL, Chang XH, Li Y, Wang Y, Guan J, et al. Effects of previous laparoscopic surgical diagnosis of endometriosis on pregnancy outcomes. *Chin Med J (Engl)* 2017;130(4):428-33. [PUBMED](#) | [CROSSREF](#)
- Saraswat L, Ayansina DT, Cooper KG, Bhattacharya S, Miligkos D, Horne AW, et al. Pregnancy outcomes in women with endometriosis: a national record linkage study. *BJOG* 2017;124(3):444-52. [PUBMED](#) | [CROSSREF](#)

17. Berlac JF, Hartwell D, Skovlund CW, Langhoff-Roos J, Lidegaard Ø. Endometriosis increases the risk of obstetrical and neonatal complications. *Acta Obstet Gynecol Scand* 2017;96(6):751-60.
[PUBMED](#) | [CROSSREF](#)
18. Kim SG, Seo HG, Kim YS. Primiparous singleton women with endometriosis have an increased risk of preterm birth: meta-analyses. *Obstet Gynecol Sci* 2017;60(3):283-8.
[PUBMED](#) | [CROSSREF](#)
19. Jeon YE, Cho S, Lee KE, Yang HI, Seo SK, Kim HY, et al. Prognostic factors for predicting spontaneous pregnancy after laparoscopic surgical treatment of endometriosis. *Korean J Obstet Gynecol* 2009;52(12):1287-95.
20. Cha J, Sun X, Dey SK. Mechanisms of implantation: strategies for successful pregnancy. *Nat Med* 2012;18(12):1754-67.
[PUBMED](#) | [CROSSREF](#)
21. Gylfason JT, Kristjansson KA, Sverrisdottir G, Jonsdottir K, Rafnsson V, Geirsson RT. Pelvic endometriosis diagnosed in an entire nation over 20 years. *Am J Epidemiol* 2010;172(3):237-43.
[PUBMED](#) | [CROSSREF](#)
22. Chae U, Min JY, Kim SH, Ihm HJ, Oh YS, Park SY, et al. Decreased progesterone receptor B/A ratio in endometrial cells by tumor necrosis factor-alpha and peritoneal fluid from patients with endometriosis. *Yonsei Med J* 2016;57(6):1468-74.
[PUBMED](#) | [CROSSREF](#)
23. Zanatta A, Pereira RM, Rocha AM, Cogliati B, Baracat EC, Taylor HS, et al. The relationship among HOXA10, estrogen receptor α , progesterone receptor, and progesterone receptor B proteins in rectosigmoid endometriosis: a tissue microarray study. *Reprod Sci* 2015;22(1):31-7.
[PUBMED](#) | [CROSSREF](#)
24. Yang SH, Kim HO, Sung NY, Ahn HS, Choo YS, Yang KM, et al. In vitro fertilization outcomes in patients with advanced endometriosis after surgical treatment. *Korean J Obstet Gynecol* 2011;54(9):517-22.
[CROSSREF](#)
25. Vercellini P, Parazzini F, Pietropaolo G, Cipriani S, Frattaruolo MP, Fedele L. Pregnancy outcome in women with peritoneal, ovarian and rectovaginal endometriosis: a retrospective cohort study. *BJOG* 2012;119(12):1538-43.
[PUBMED](#) | [CROSSREF](#)
26. Kortelahti M, Anttila MA, Hippeläinen MI, Heinonen ST. Obstetric outcome in women with endometriosis--a matched case-control study. *Gynecol Obstet Invest* 2003;56(4):207-12.
[PUBMED](#) | [CROSSREF](#)
27. Kuivasaari-Pirinen P, Raatikainen K, Hippeläinen M, Heinonen S. Adverse outcomes of IVF/ICSI pregnancies vary depending on aetiology of infertility. *ISRN Obstet Gynecol* 2012;2012:451915.
[PUBMED](#) | [CROSSREF](#)
28. Smithers PR, Halliday J, Hale L, Talbot JM, Breheny S, Healy D. High frequency of cesarean section, antepartum hemorrhage, placenta previa, and preterm delivery in in-vitro fertilization twin pregnancies. *Fertil Steril* 2003;80(3):666-8.
[PUBMED](#) | [CROSSREF](#)
29. Haas D, Wurm P, Shamiyeh A, Shebl O, Chvatal R, Oppelt P. Efficacy of the revised Enzian classification: a retrospective analysis. Does the revised Enzian classification solve the problem of duplicate classification in rASRM and Enzian? *Arch Gynecol Obstet* 2013;287(5):941-5.
[PUBMED](#) | [CROSSREF](#)
30. Conti N, Cevenini G, Vannuccini S, Orlandini C, Valensise H, Gervasi MT, et al. Women with endometriosis at first pregnancy have an increased risk of adverse obstetric outcome. *J Matern Fetal Neonatal Med* 2015;28(15):1795-8.
[PUBMED](#) | [CROSSREF](#)