

Usefulness of intraductal RFA in patients with malignant biliary obstruction

Sung Yong Han, MD^a, Dong Uk Kim, MD, PhD^{b,*}, Dae Hwan Kang, MD, PhD^c, Dong Hoon Baek, MD, PhD^b, Tae Hoon Lee, MD, PhD^d, Jae Hee Cho, MD, PhD^{e,*}

Abstract

Background/Aims: Intraductal radiofrequency ablation (ID-RFA) is a novel therapy for unresectable malignant biliary obstructions. ID-RFA for perihilar lesions is associated with a high risk of adverse events. We aimed to evaluate the feasibility and efficacy of temperature-controlled ID-RFA for perihilar malignant biliary obstruction.

Methods: Sixteen patients with pathologically proven perihilar cholangiocarcinoma were prospectively enrolled to evaluate the feasibility of hilar ID-RFA. Clinical efficacy and outcomes were subsequently evaluated in a multicenter retrospective cohort.

Results: Nine of the 16 patients in the prospective cohort had Bismuth type IV and 7 had type IIIA perihilar cholangiocarcinoma. The median length of stricture was 34.5 mm. The median number of ID-RFA sessions was three, and all sessions were technically and functionally successful without severe adverse events. Clinical outcomes were assessed using a multicenter hilar ID-RFA cohort of 21 patients; the median stent patency and overall survival were 90 days (range: 35–483 days) and 147 days (range: 92–487 days), respectively. An approximate 16-month patency of the bile duct was maintained in one patient who had an intraductal growth pattern. In a comparison of the self-expandable metallic stent (SEMS) and plastic stent (PS) after hilar ID-RFA, no differences in stent patency (89 vs 90.5 days, respectively; $P = .912$) and adverse events (20.0% vs 10%, respectively; $P = .739$) were observed.

Conclusions: ID-RFA at 7 W for 120 seconds is safe and feasible in patients with advanced perihilar cholangiocarcinoma. After ID-RFA, SEMS and PS placement showed comparable patency and survival rates.

Trial registration number: KCT0003223

Abbreviations: ERCP = endoscopic retrograde cholangiopancreatography, ID-RFA = Intraductal radiofrequency ablation, PS = plastic stent, PTBD = percutaneous transhepatic biliary drainage, RFA = radiofrequency ablation, SEMS = self-expandable metallic stent.

Keywords: biliary tract cancer, ERCP, radiofrequency ablation, perihilar cholangiocarcinoma, self-expandable metallic stents

1. Introduction

Local ablation therapy is performed to maintain ductal patency, which improves survival and quality of life in patients with unresectable hilar cholangiocarcinoma. The endoscopic deployment of a self-expandable metallic stent (SEMS) is the standard palliation in these patients, and a SEMS has a life expectancy over

3 months, providing a high efficacy of bile drainage, low incidence of complications, and patient acceptance.^[1] However, the efficient drainage of hilar biliary strictures is challenging because of long and tight strictures, acute angulation of the bile duct, and complicated structures, such as the portal vein. The successful biliary drainage remains controversial especially as

This study was registered at the Clinical Research Information Service (KCT 0003223).

This study was supported by a grant from the National R&D Program for Cancer Control, Ministry of Health and Welfare, Republic of Korea (HA20C0009).

The authors have no conflicts of interest to disclose.

All data generated or analyzed during this study are included in this published article [and its supplementary information files].

^a Department of Internal Medicine and Biomedical Research Institute, Pusan National University Hospital, Busan, ^b Department of Internal Medicine and Biomedical Research Institute, Pusan National University Hospital, Pusan National University School of Medicine, Busan, South Korea, ^c Department of Internal Medicine, Medical Research Institute, Pusan National University School of Medicine and Research Institute for Convergence of Biomedical Science and Technology, Pusan National University Yangsan Hospital, Yangsan-si, Gyengsangnam-do, ^d Digestive Disease Center and Research Institute, Department of Internal Medicine, SoonChun-Hyang University School of Medicine, Cheonan, ^e Department of Internal Medicine, Gangnam Severance Hospital, Yonsei University College of Medicine, Seoul, South Korea.

* Correspondence: Dong Uk Kim, Division of Gastroenterology, Department of Internal Medicine and Biomedical Research Institute, Pusan National University Hospital, Gudeok ro 179, Seo-gu, Busan, Republic of Korea 49241 (e-mail: amlm3@hanmail.net); Jae Hee Cho, Department of Internal Medicine, Gangnam Severance Hospital, Yonsei University College of Medicine, 211, Eonju-ro Gangnam-gu, Seoul 06273, Republic of Korea (e-mail: jhcho9328@gmail.com).

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Han SY, Kim DU, Kang DH, Baek DH, Lee TH, Cho JH. Usefulness of intraductal RFA in patients with malignant biliary obstruction. *Medicine* 2020;99:33(e21724).

Received: 11 July 2020 / Accepted: 13 July 2020

<http://dx.doi.org/10.1097/MD.00000000000021724>

regards the use of a SEMS or plastic stent (PS). Both stent types completely occlude within 6 months despite improvements in anticancer treatments for perihilar cholangiocarcinoma; therefore, the development of a novel treatment method to prolong stent patency is necessary.

Newer endoscopic palliative therapies, including local ablative techniques, have been introduced. Photodynamic therapy has been reported to be an effective adjunctive therapy in unresectable cholangiocarcinoma.^[2] However, photodynamic therapy requires specialized facilities, has uncertain survival benefits, and can have complications, such as photosensitivity.^[3] Intraductal radiofrequency ablation (ID-RFA) using an endobiliary radiofrequency ablation (RFA) catheter has been identified as another treatment option. ID-RFA is easily performed on distal malignant biliary obstructions, and the local tumor is successfully ablated with minimal complications. However, ID-RFA in perihilar applications has the potential to cause bile duct perforation and vascular injuries, such as hemobilia and liver infarction, because the hilar bile duct is located near the hepatic artery and portal vein complexes.^[4–6] We aimed primarily to evaluate the feasibility and clinical outcomes of temperature-controlled hilar ID-RFA in patients with perihilar malignant biliary obstruction. Secondly, we sought to compare the clinical efficacy of a bilateral SEMS and PS after hilar ID-RFA.

2. Materials and methods

2.1. Study design and population

We performed a prospective, uncontrolled, single-center study that included 18 patients with unresectable perihilar cholangiocarcinoma. Two patients had a choledochoduodenostomy and were excluded from the analysis. The inclusion criterion was pathologically confirmed perihilar cholangiocarcinoma of Bismuth type III or IV. The decision of whether a patient had an unresectable malignancy was based on their medical fitness or disease extent (tumor extending into both liver lobes, major vascular involvement, or Bismuth type IV). The ethics committee of the institutional review board of the hospital approved this prospective study (H-1401-005-013), which was registered in the Clinical Research Information Service (KCT0003223). After a feasibility evaluation, a further retrospective analysis of clinical outcomes regarding hilar ID-RFA was conducted using a multicenter hilar ID-RFA cohort of 21 patients (an additional 5 patients underwent hilar ID-RFA at 7 W and 120 seconds from Gil Medical Center) (Fig. 1). The baseline characteristics of

patients from each hospital were not statistically different between the two groups (data not shown).

2.2. Clinical trial protocol: ID-RFA procedure

All patients underwent endoscopic retrograde cholangiopancreatography (ERCP) before enrollment for tissue confirmation and biliary drainage. ID-RFA was subsequently performed using a newly designed ID-RFA catheter (ELRA STARmed, Seoul, Korea) and a transpapillary approach. The ID-RFA catheter is a single-use, disposable, bipolar device that is suitable for endoluminal delivery of RFA into the biliary tree over a guidewire. In addition, the catheter can be deployed through a side-view endoscope with 4.2-mm diameter working channels (Olympus TJF-260 V, Tokyo, Japan) or a forward-view endoscope with 3.8-mm diameter working channels (Olympus GIF-260, Tokyo, Japan). Each ID-RFA session was applied at 7 W (11 or 18 mm) for 120 s under an 80°C intraductal target temperature. During the procedure, the wattage was automatically guided by a STARmed RF generator (STARmed, Seoul, Korea). The prevention of tissue charring is the major advantage of temperature-controlled ID-RFA; this may reduce the risk of ID-RFA induced perforation and bleeding. The actual ID-RFA procedure is as follows: The ID-RFA catheter is placed within the tumor stricture under fluoroscopic visualization of the biliary system, and repeated applications of ID-RFA are carefully made by passing the short-length ID-RFA catheter (11 or 18 mm) to the treated segments (1–5 applications per intervention); after ID-RFA treatment, either a bilateral Y-configured SEMS (Niti-S Biliary Y or T-stent, Taewoong, Goyang, Korea; Niti-S or Niti-D stent, Taewoong Medical, Goyang, Korea), bilateral uncovered SEMS (Benefit stent, MiTec, Seoul, Korea), or bilateral PS (Boston scientific, Boston, MA, USA & Cook medical, Bloomington, IN, USA) is placed depending on the clinical condition of each patient.

2.3. Outcome measurement parameters

We evaluated adverse events and biliary patency. The outcome of bilateral drainage was evaluated according to the following parameters:

- 1) Technical success was defined as the proper placement of the ID-RFA catheter in the stricture site and successful ablation, including a successful bilateral SEMS or PS placement across the stricture.
- 2) Functional success was defined as radiographically adequate positioning, immediate biliary decompression, and at least a 30% reduction of serum bilirubin level within 2 weeks.
- 3) Procedure-related adverse events and mortality.
- 4) Stent patency (the period between stent insertion and occlusion or patient's death). Occlusion of the stent was confirmed if patients had recurrent or persistent jaundice with cholestasis or cholangitis (leukocytosis, fever, increasing bilirubin).

2.4. Statistical analysis

Statistical analysis was performed using SPSS software (version 22.0, IBM Corp., Armonk, NY). Categorical data were expressed as frequencies and percentages, and between-group differences were evaluated using the Fisher Exact test. Continuous data were




Figure 1. Flowchart of the study.

expressed as medians with ranges, and between-group differences were evaluated using the Mann-Whitney *U* test. Statistical significance was set at $P < .05$. For comparison, stent patency was assessed using Kaplan–Meier survival curves and the log-rank test.

3. Results

3.1. Patient characteristics and clinical outcomes of ID-RFA in the prospective feasibility cohort

Figure 1 shows the flowchart of this study. Detailed information on the patient characteristics and results of the ID-RFA procedure in the prospective cohort is shown in Table 1. The 16 enrolled patients included 6 women (median age: 75.5 years; range: 59–85 years). Ten patients had hypertension, 4 patients had diabetes mellitus, and no patients had liver cirrhosis or viral hepatitis. Of the 16 enrolled patients, 9 had Bismuth type IV and 7 had type IIIA perihilar cholangiocarcinoma. The median stricture length was 34.5 mm (range: 17–63 mm). The median number of ID-RFA applications per single session was 3 (range: 1–5). ID-RFA was performed in 10 patients using an 11-mm RFA catheter (62.5%), in 7 patients using an 18-mm catheter, and in 1 patient using both catheters.

The clinical outcomes of the ID-RFA procedure in the prospective cohort are shown in Table 2. The median initial total bilirubin level was 10.39 mg/dL. In all patients, ID-RFA was technically and functionally successful. After ID-RFA, 6 patients had placement of a bilateral SEMS, 9 patients had placement of a bilateral PS, and 1 patient had insertion of an endoscopic nasobiliary drainage (ENBD) tube. The median total bilirubin level was 2.18 mg/dL at 1 day after ID-RFA and 1.30 mg/dL at 2 weeks after ID-RFA. No fatal adverse events, such as major bleeding or perforation, were observed. The median duration of stent patency was 91.5 days (35–483 days). The median overall survival period was 131 days (50–487 days). No difference in stent patency between the bilateral SEMS and PS groups (93.5 vs 91 days, respectively, $P = .607$) was observed. Interestingly, 1 patient who had an intraductal growth pattern was eligible for repeated ID-RFA and survived for 332 days without internal or external biliary drainage.

Table 1
Baseline characteristics and result of ID-RFA procedure.

Variables	Patients (n = 16)
Age, median (range)	75.5 (59–85)
Sex, Male, n (%)	10 (62.5)
Bismuth type IIIA/IV, n (%)	9/7 (56.3/43.8)
Liver cirrhosis, n (%)	0 (0)
Viral hepatitis, n (%)	0 (0)
Stricture length, mm, median (range)	34.5 (17–63)
ID-RFA probe length, n (%)	
11 mm/18 mm*	10/7 (62.5/43.8)
Ablation power, n (%)	
7 W	16 (100)
Ablation duration, n (%)	
120 s	16 (100)
Bilateral stenting, n (%)	15 (93.8)
Number of ablation attempts in one session, median (range)	3 (1–5)

* one patient was used 2 type of ID-RFA probe.
ID-RFA = intraductal radiofrequency ablation.

3.2. Clinical outcomes in the multicenter hilar ID-RFA cohort

Table 3 shows the clinical outcomes and baseline characteristics of patients in the multicenter hilar RFA cohort. The baseline characteristics did not differ significantly between the SEMS and PS groups. The median stricture length was 34 mm (range: 17–63 mm). The median number of ID-RFA applications per single session was 3 (range: 1–5). ID-RFA was performed in 15 patients using an 11-mm RFA catheter (71.4%), in 7 patients using an 18-mm catheter, and in 1 patient using both catheters. The median duration of stent patency was 90 days (35–483 days), and no difference was observed in the stent patency between the SEMS and PS groups following ID-RFA (median stent patency: 89 vs 90.5 days, respectively; $P = .912$). The functional duration of the stent was not successfully identified in 2 patients; therefore, the last follow-up visit was considered as the stent occlusion. The median overall survival period was 147 days (50–487 days); the SEMS group showed slightly longer overall survival than the PS group, but the difference was not statistically significant (median overall survival: 157 vs 119.5 days; $P = .142$). Only 2 patients in the SEMS group received chemotherapy.

One patient with cholangitis was controlled with intravenous antibiotics, and 2 patients with cholecystitis underwent percutaneous cholecystostomy. In both the SEMS and PS groups, no significant difference in the adverse event rate was observed (20% vs 10%, respectively, $P = .739$). Interestingly, the ENBD tube was removed in one patient with an intraductal growth pattern, and drainage was possible for a long period without any additional procedure (483 days). Figure 2 shows the result of stent patency according to intraductal growth pattern, and no significant difference between the two groups was observed. However, the patients with an intraductal growth pattern tended to have longer stent patency than other patients (106 vs 89 days, $P = .097$).

4. Discussion

Temperature-controlled ID-RFA was safe and feasible in patients with advanced hilar cholangiocarcinoma and had no major adverse events, such as perforation and hemobilia. For the hilar ID-RFA, we used a short ID-RFA catheter under a low power (7W) for a duration of 120 seconds. Because stricture length has a wide range (17–63 mm), we performed multiple overlapping ID-RFAs to cover the entirety of perihilar strictures. RFA can also be applied to more than one stricture site during the same session, such as the common hepatic duct or left- or right-hepatic duct, in patients with Klatskin tumors. In the prospective feasibility study, technical and clinical success was achieved in all enrolled patients, and our adverse event rate of 18.8% is comparable to that of other studies (0–33.3%).^{17–10}

No specific recommendation has been given for the ID-RFA settings of duration, energy, and technique for hilar applications. The actual ID-RFA ablation depth in the human bile duct is approximately 4 mm (1.7–4.3 mm), and the effective length of bile duct ablation has been estimated to be 40% to 72% of the length of the RFA catheter.^{11–13} A previous in vivo swine study showed that hilar ID-RFA related perforation in the normal bile duct occurred when a duration of over 90 seconds was used.¹⁷ However, some clinical studies of hilar ID-RFA revealed no perforation at a duration up to 120 seconds.^{9,11} Since a cancerous bile duct has a thicker wall than a normal bile duct, we chose to perform hilar ID-RFA for a duration up to 2 minutes to get better outcomes. We chose shorter ID-RFA catheters because longer catheters are not suitable for

Downloaded from http://journals.lww.com/md-journal by BNDMfsePHKav1ZEoum1tQINa+kLNEzjgsho4XM0hOy WCX1AWnYQpII0HHD33DDO0dFy7TVSfACf3V/C1y0abggQZXdwmfKZBYtws= on 08/09/2023

Table 2
Clinical outcomes of ID-RFA in prospective feasibility cohort.

Variables	Patients (n=16)	bilateral SEMS (n=6)	bilateral PS (n=9)	P value
Technical success, n (%)	16 (100)	6 (100)	9 (100)	>.99
Clinical success, n (%)	16 (100)	6 (100)	9 (100)	>.99
Total bilirubin, median (range), mg/dL				
At admission	10.39 (0.25–24.8)	8.18 (1.4–24.8)	11.09 (0.25–19.31)	.776
Pre- ID-RFA (post ERCP)	2.22 (0.27–9.07)	1.22 (0.66–3.79)	2.8 (0.27–9.07)	.272
1 day after ID-RFA	2.18 (0.42–11.81)	1.53 (0.86–3.83)	2.65 (0.42–7.35)	.272
2 weeks after ID-RFA	1.30 (0.55–4.58)	0.83 (0.60–2.10)	1.45 (0.55–4.58)	.088
Adverse events, n(%)	3 (18.8)	2 (33.4)	1 (11.1)	.529
Perforation	0 (0)	0 (0)	0 (0)	
Hemobilia	0 (0)	0 (0)	0 (0)	
Cholangitis	1 (6.3)	1 (16.7)	0 (0)	.205
Cholecystitis	2 (12.5)	1 (16.7)	1 (11.1)	.864
pancreatitis	0 (0)	0 (0)	0 (0)	
Others	0 (0)	0 (0)	0 (0)	
Chemotherapy	2 (12.5)	2 (33.3)	0 (0.0)	.205
Stent patency duration, median (range), d	91.5 (35–483)	93.5 (50–172)	91 (35–133)	.607
30-day stent patency, n (%)	16 (100)	6 (100)	9 (100)	>.99
90-day stent patency, n (%)*	10/14 (71.4)	3/4 (75.0)	6/9 (66.7)	.764
Survival days, median (range), d	131 (50–487)	133 (50–302)	129 (63–168)	.689

SEMS = self expandable metallic stent, PS = plastic stent.

* Two patients were lost to follow up.

intraductal ablation, since the perihilar bile duct is angulated and separated, and longer catheters have the potential to cause biliovascular complications. Therefore, we recommended the use of low power and short ID-RFA catheters (7W and 11 mm or 18 mm) for the safe and successful performance of hilar ID-RFA. When dealing with long segments of bile duct cancer, repeated ID-RFA with a

tandem overlapping of RFA catheters can overcome the limitations of ID-RFA.^[7] Furthermore, since ERCP is not always successful when a perihilar biliary obstruction is present, the percutaneous transhepatic biliary drainage (PTBD) tract can be used as an alternative to ERCP-guided RFA.^[7–10] Consequently, the utilization of a multidisciplinary care approach through the effective applica-

Table 3
Clinical outcome of multicenter hilar RFA cohort.

Variables	Patients (n=21)	bilateral SEMS (n=10)	bilateral PS (n=10)	P value
Age, median (range)	76 (59–85)	76 (49–81)	76.5 (65–85)	.529
Sex, Male, n (%)	14 (66.7)	6 (60.0)	7 (70.0)	.739
Bismuth type IIIA/IV (%)	8/13 (38.1/61.9)	2/8 (20.0/80.0)	5/5 (50.0/50.0)	.280
Stricture length, mm, median (range)	34 (17–63)	32.5 (20–59)	35.5 (17–63)	.912
ID-RFA probe length, n (%)				
11 mm/18 mm*	15/7 (71.4/33.3)	8/3 (80.0/30.0)	6/4 (60.0/40.0)	.853
Number of ablation attempts in one session, median (range) †	3 (1–5)	3 (2–4)	3 (1–5)	.796
Total bilirubin, median (range), mg/dL				
At admission	9.87 (0.25–24.8)	10.80 (1.4–24.8)	10.39 (0.25–19.31)	>.99
pre- ID-RFA (post ERCP)	2.62 (0.27–9.07)	1.58 (0.66–7.30)	2.71 (0.27–9.07)	.853
1 day after ID-RFA	2.28 (0.42–11.81)	1.82 (0.86–5.8)	2.47 (0.42–7.35)	.912
2 weeks after ID-RFA	1.36 (0.55–4.58)	1.02 (0.60–2.10)	1.41 (0.55–4.58)	.315
Adverse events, n(%)	3 (14.3)	2 (20.0)	1 (10.0)	.739
perforation	0 (0)	0 (0)	0 (0)	
Hemobilia	0 (0)	0 (0)	0 (0)	
Cholangitis	1 (4.8)	1 (10.0)	0 (0)	.739
Cholecystitis	2 (9.5)	1 (10.0)	1 (10.0)	>.99
Pancreatitis	0 (0)	0 (0)	0 (0)	
Others	0 (0)	0 (0)	0 (0)	
Chemotherapy	2 (9.5)	2 (20.0)	0 (0.0)	.739
Stent patency duration, median (range), d	90 (35–483)	89 (50–172)	90.5 (35–133)	.912
30-day stent patency, n (%)	21 (100)	10 (100)	10 (100)	>.99
90-day stent patency, n (%)‡	11/19 (57.9)	4/8 (50.0)	6/10 (60.0)	.762
Survival days, median (range), d	147 (50–487)	157 (50–302)	119.5 (63–168)	.143

SEMS = self expandable metallic stent, PS = plastic stent.

* one patient was used two type of ID-RFA probe.

† ID-RFA was performed by 7 watts for 120 s.

‡ Two patients were lost to follow up.

Downloaded from http://journals.lww.com/rnd-journal by BMDM5ePHKav17Eoum1t1QINa+kLLNEZgshHo4XMM10iOy WOX1AWmYQpII0HrHD333DOODfyt7TVSFACf3V/C1y0abgqZxdtwIKZBYtws= on 08/09/2023




Figure 2. Kaplan-Meier curve for stent patency according to whether or not of intraductal growth pattern.

tion of ERCP and PTBD is necessary in patients with a perihilar biliary obstruction.

The multicenter hilar ID-RFA cohort had a median stent patency of 90 days, and survival was 147 days; these results are similar to those of a Klatskin tumor without hilar ID-RFA. Therefore, we could not confirm whether the hilar RFA was effective for these patients. Most patients in this study were in poor health, which made additional chemotherapy impossible. Therefore, we admit that the actual efficacy of hilar ID-RFA could not be verified in this study. In addition, we compared the clinical usefulness of a PS to a SEMS after hilar ID-RFA because insufficient data is available on which stent to use after hilar ID-RFA. Our study shows no difference in stent patency, survival, and adverse events between a PS and SEMS. In our study, a SEMS did not show a superior effect compared to a PS in hilar malignancy, unlike other studies, and the replaceable PS allows for repeated RFA, which is associated with a survival benefit.^[14] A PS may be a more reasonable therapeutic option after hilar ID-RFA; however, this should be confirmed through further studies because the selection of stent type depends on the patient's condition.

Determining a subgroup that can expect a therapeutic effect after hilar ID-RFA is very important when deciding whether to perform hilar ID-RFA. Notably, an approximate one-year patency of the bile duct was achieved in one patient who had an intraductal growth pattern of bile duct cancer; this patient did not receive any additional therapy, including chemotherapy. The intraductal growth pattern cholangiocarcinoma constitutes 8% to 18% of cholangiocarcinomas.^[15-17] In our study, a total of 4 patients had cholangiocarcinoma with an intraductal growth pattern. We observed a tendency to maintain longer stent patency in the patients with an intraductal growth pattern than that in the patients without an intraductal growth pattern, but the difference was not statistically significant. Therefore, cholangiocarcinoma with an intraductal growth pattern may show a better response to

RFA treatment. One patient was able to maintain stent patency for 1 year through repeated ID-RFA; such repeated RFA is thought to help extend the total duration of stent patency. Further research is required to determine whether patients with repeated RFA and an intraductal growth pattern have a good prognosis.

Our research has several limitations. First, we enrolled a small number of patients who were treated in a single institution in the prospective feasibility cohort and did not have a control group. Further randomized, controlled studies are needed to prove the usefulness of ID-RFA for perihilar cholangiocarcinoma. Second, patients with advanced diseases were predominantly enrolled, and the proportion of patients treated with chemotherapy was small. In particular, the advanced stages were analyzed heterogeneously, including locally advanced stages and distant metastases. Furthermore, the SEMS group had a heterogeneous mix of procedures, such as stent-in-stent or stent-by-stent, and various stent diameters were used (8mm/10mm). Finally, the proportion of patients for whom follow-up was discontinued was high (33.3%), and this affected the accuracy of the assessment.

In conclusion, ID-RFA at 7W for 120 seconds is safe and feasible for advanced hilar obstruction. After ID-RFA, both SEMS and PS placement showed comparable patency and survival rates. Cholangiocarcinoma with an intraductal growth type and repeated ID-RFA is thought to be a way to maximize the effect of ID-RFA. For repeated ID-RFA, a PS after ID-RFA may be beneficial. Long-term survival gains could be achieved if these local therapies are used appropriately. However, randomized, controlled studies and multi-institutional studies, especially ones that compare RFA with a control, need to be conducted to confirm the benefits of long-term stent patency and patient survival.

Author Contribution

Critical revision: Jae Hee Cho, Tae Hoon Lee.

Design of the work: Dong Uk Kim, Dae Hwan Kang.

Drafting the article: Sung Yong Han.

Data acquisition & analysis: Sung Yong Han, Dong Hoon Baek.

References

- Rustagi T, Jamidar PA. Intraductal radiofrequency ablation for management of malignant biliary obstruction. *Dig Dis Sci* 2014;59:2635-41.
- Moole H, Tathireddy H, Dharmapuri S, et al. Success of photodynamic therapy in palliating patients with nonresectable cholangiocarcinoma: a systematic review and meta-analysis. *World J Gastroenterol* 2017;23:1278.
- Pereira SP, Jitlal M, Duggan M, et al. PHOTOSTENT-02: porfimer sodium photodynamic therapy plus stenting versus stenting alone in patients with locally advanced or metastatic biliary tract cancer. *ESMO Open* 2018;3:e000379.
- Zhou C, Wei B, Gao K, et al. Biliary tract perforation following percutaneous endobiliary radiofrequency ablation: a report of two cases. *Oncol Lett* 2016;11:3813-6.
- Topazian M, Levy M, Patel S, et al. Hepatic artery pseudoaneurysm formation following intraductal biliary radiofrequency ablation. *Endoscopy* 2013;45:E161-2.
- Alis H, Sengoz C, Gonenc M, et al. Endobiliary radiofrequency ablation for malignant biliary obstruction. *Hepatobiliary Pancreat Dis Int* 2013;12:423-7.
- Kim EJ, Cho JH, Kim YJ, et al. Intraductal temperature-controlled radiofrequency ablation in malignant hilar obstruction: a preliminary study in animals and initial human experience. *Endosc Int Open* 2019;7:E1293-300.

- [8] Cui W, Wang Y, Fan W, et al. Comparison of intraluminal radiofrequency ablation and stents v. stents alone in the management of malignant biliary obstruction. *Int J Hyperthermia* 2017;33:853–61.
- [9] Jeon CH, Kim CW, Kwon H. Safety and efficacy of percutaneous intraductal radiofrequency ablation followed by dual stent placement for malignant biliary hilar obstruction. *J Kor Soc Radiol* 2017;77:298–308.
- [10] Nayar MK, Oppong KW, Bekkali NL, et al. Novel temperature-controlled RFA probe for treatment of blocked metal biliary stents in patients with pancreaticobiliary cancers: initial experience. *Endosc Int Open* 2018;6:E513–7.
- [11] Kim EJ, Chung DH, Kim YJ, et al. Endobiliary radiofrequency ablation for distal extrahepatic cholangiocarcinoma: a clinicopathological study. *PloS One* 2018;13:e0206694.
- [12] Itoi T, Isayama H, Sofuni A, et al. Evaluation of effects of a novel endoscopically applied radiofrequency ablation biliary catheter using an ex-vivo pig liver. *J Hepatobiliary Pancreat Sci* 2012;19:543–7.
- [13] Cho JH, Lee KH, Kim JM, et al. Safety and effectiveness of endobiliary radiofrequency ablation according to the different power and target temperature in a swine model. *J Gastroenterol Hepatol* 2017;32:521–6.
- [14] Yang J, Wang J, Zhou H, et al. Efficacy and safety of endoscopic radiofrequency ablation for unresectable extrahepatic cholangiocarcinoma: a randomized trial. *Endoscopy* 2018;50:751–60.
- [15] Seo N, Kim DY, Choi J-Y. Cross-sectional imaging of intrahepatic cholangiocarcinoma: development, growth, spread, and prognosis. *Am J Roentgenol* 2017;209:W64–75.
- [16] Yamasaki S. Intrahepatic cholangiocarcinoma: macroscopic type and stage classification. *J Hepatobiliary Pancreat Surg* 2003;10:288–91.
- [17] Lee JW, Han JK, Kim TK, et al. CT features of intraductal intrahepatic cholangiocarcinoma. *Am J Roentgenol* 2000;175:721–5.